Effect of Vitomolt supplements in feed on growth and survival rate of white shrimp (Litopenaeus vannamei) seeds.

by Dr. Ir. Hadijah, Msi

Submission date: 07-Feb-2021 02:22PM (UTC+0700)
Submission ID: 1503447138
File name: ijsrp-p10778.pdf (341.88K)
Word count: 2966
Character count: 15291

Effect of Vitomolt supplements in feed on growth and survival rate of white shrimp (*Litopenaeus vannamei*) seeds.

Zainuddin[1]*, Y. Fujaya1, M.I. Djawad[1], S. Aslamyah[1], K. Nur1, Hadijah[2]

[1] Fisheries Department, Faculty of Marine and Fisheries Science, Universitas Hasanuddin, Makassar
[2] Aquaculture Study Program, Faculty of Agriculture, Universitas Bosowa, Makassar

DOI: 10.29322/IJSRP.10.11.2020.p10778
http://dx.doi.org/10.29322/IJSRP.10.11.2020.p10778

Abstract- Feed is a very important factor in vaname shrimp aquaculture. The addition of phytoecdysteroid supplements (vitomolts) to the feed can increase protein retention, thus the shrimp growth will be better. This study aimed to determine the appropriate dose of Vitomolt supplement in feed for growth and survival of vaname shrimp seeds. The results of this study are expected to be useful in determining the best Vitomolt supplementation in the vaname shrimp feed formulation. The experimental animal used was the vaname shrimp seeds with an average initial weight of 1.5 g per individual. The research design used was a completely randomized design (CRD) with 4 treatments, namely treatment A without Vitomolt supplement, B 1.5 g Vitomolt /kg feed, C 3.0 g Vitomolt/kg feed, and D 4.5 g Vitomolt /kg feed. Each treatment had 3 replications, hence there were 12 experimental units. The experimental feed application was carried out for 40 days of maintenance. The results showed that the treatment of Vitomolt supplement dosage in the feed had a significant effect on the daily growth rate and survival rate of vannamei shrimp. The results of further tests showed that the vitomolt dose of 4.5 g/kg of feed was significantly different from the treatment without vitomolt, but did not differ from those with 1.5 g and 3.0 g /kg of feed Vitomolt dose on the daily growth rate of vaname shrimp seeds. Furthermore, the Vitomolt dose of 3.0 g /kg of feed was significantly different from the treatment without Vitomolt but not significantly different from 1.5 g and 4.5 g /kg of feed Vitomolt dose, on the survival rate of the vaname shrimp seeds. Based on the results of this study, it is recommended to add 3.0 g /kg of vitomolt for vaname seeds feed.

Index Terms- feed, growth, survival rate, Vitomolt

I. Introduction

In an intensive vaname shrimp culture system in ponds, feed is one of the strategic components that determine the success of the business. In these activities, nearly 60-70% of the total production costs are used for the purchase of feed (Haryati et al. 2009; Haliman and Dian, 2005). The high price of commercial shrimp feed is due to the fact that almost 100% of the fish meal used as a protein raw material is imported from other countries. Therefore, the amount of protein in the feed must be limited, protein is optimized only for growth, while energy needs are met from carbohydrates (*protein-sparring effect by carbohydrates*) which are cheaper (Rosas et al. 2001). However, the problem is when the protein content of the feed is low, the potential for protein retention by cultured organisms is also low. It is in this condition that the vitomolts as phytoecdysteroids play an important role in increasing protein retention (Klein 2004, Fujaya et al. 2011).

According to Aslamyah (2000), one of the efforts to reduce protein composition in feed, without disturbing the growth of cultivated organisms is to use steroid hormones. This occurs, because steroid hormones are receptors that carry protein into cells, so that they can activate protein metabolism. Spinach extract, which is a phytoecdysteroid including the steroid class, when added to feed, in addition to accelerating molting and growth, is also expected to increase the efficiency of feed protein utilization. The results showed that vitomolts can retain protein up to> 40%. Thus, vitomolts not only have the potential to reduce feed prices and aquaculture management costs, but also play a direct role in increasing production. Experience with crab can be applied to shrimp with a few adjustments. Because the growth of crab and shrimp are both controlled by the molting hormone in the form of ecdysteroid.

The results of research by Wahyuningsih (2008) and Susanti (2009) suggest that the right dose of ecdysteroid will stimulate crustaceans to molt optimally, while low doses cannot stimulate molting, otherwise too high doses cause inhibition. Molting is an important initial phase for crabs and shrimp because by molting they can experience growth. This indicates that the ecdysteroid dose plays an important role in vitomolt supplementation in the feed. Until now, the dose of vitomolts as a supplement in vaname shrimp feed has never been studied, so it is necessary to do research on this matter. It is hoped that this research will produce low protein vaname shrimp feed with vitomolt supplements as ecdysteroid so that shrimp growth can be faster and the impact of nitrogen waste into the environment is minimal.

II. Methodology

The research was conducted at the Mini Hatchery, Faculty of Marine Sciences and Fisheries Hasanuddin University. The containers used in this study were 12 glass aquariums with a

This publication is licensed under Creative Commons Attribution CC BY.

capacity of 72 liters and filled with 56 L of brackish water for each aquarium. The water used is sea water that has been diluted to reach a salinity of 25-27 ppt. The test animal used in this study was juvenile shrimp with a weight of 1.5 g per individual. Each container is scattered with 20 vaname shrimps. Before giving the test feed, the shrimp were adapted to commercial feed for a week. This is so that the shrimp gets used to eating pellets. After the adaptation period is complete, the test shrimp are fed according to the treatment.

This study used a formulated feed with a low protein composition. The feed raw materials for the formulations are sourced from fish meal, soybean flour, corn flour, sweet potato flour, CMC, vitamin, mineral mix and fish oil as well as vitomolts as a supplement. Feed making begins by pulverizing all the dry ingredients used. All materials are weighed as needed and placed in plastic bags.

All dry feed ingredients are mixed starting from small amounts of fine ingredients followed by large amounts of raw materials, then stirring until they are well blended. Furthermore, added fish oil, vitamin and mineral mix. After it is evenly mixed, 400 mL of warm water is added to the mixture of 1 kg of feed raw materials to form a dough/paste. The feed dough is stirred until it doesn't stick to the hands. Then the dough is put into a feed molding device and molded into pellets. The pelleted feed is spread regularly on trays and dried in the sun to dry (Zainuddin et al. 2019). The dry pelleted feed is mixed with vitomolts according to the treatment dose. The feed that has been mixed with vitomolt is air dried and stored in a plastic jar and labeled according to treatment.

The formulated feed which was tested according to the treatment was randomly placed into the seed rearing medium. The amount of feed given is 10% of the body weight of the seeds (Zainuddin et al. 2020). Feeding is done with a frequency of four times every day, namely 06.00 AM, 10.00 AM, 14.00 PM and 18.00 PM. The test feed application was carried out for 40 days of maintenance to see the response to be measured.

Shrimp growth data were taken per week by sampling shrimp in each aquarium. Shrimp weight was weighed using a digital scale with a capacity of 500 g. Growth sampling was carried out every week to determine the increase in body weight of shrimp and to adjust feed. The method used is an experimental method by applying vitomolt supplementation at various doses as a treatment. The experimental design used was a completely randomized design (CRD) with 4 treatments, namely A without vitomolt supplementation, B 1.5 g vitomolt / kg feed, C 3.0 g vitomolt / kg feed, D 4.5 g vitomolt / kg feed. Each treatment had 3 replications in order to obtain 12 experimental units.

The measured research parameters measured in this study are:
1. Survival Rate (SR)
SR = Nt / Nox100
where :
Nt = number of individuals at the end of the study (individu)
No = number of individuals at the beginningt of the study (individu)
2. Specific growth rate (SGR)

$$SGR = (Ln\ Wt - Ln\ Wo) / t \times 100$$

Where:
Wt = average individual weight at the end of the study (g)
Wo = average individual weight at the beginning of the study (g)
t = maintenance time (days)

Data were analyzed using analysis of variance (ANOVA) using SPSS version 24 software.

III. Results and Discussion

Daily Growth Rate

The average daily growth rate of vaname shrimp at the end of the study is presented in Table 1.

Table 1. Daily growth rate (DGR) of vaname shrimp during the study

Vitomolt dose (g/kg of feed)	DGR (%/day)
(A) Control	2.11 ± 0.29^{a}
(B) 1.5	2.39 ± 0.41^{ab}
(C) 3.0	3.04 ± 0.51^{ab}
(D) 4.5	3.38 ± 0.32^{b}

Note: Different letters in the DGR column indicate significant differences between treatments (P <0.05)

The results of analysis of variance (ANOVA) showed that the addition of vitomolt to pelleted feed had a significant effect (p <0.05) on the daily growth rate of vannamei shrimp. Furthermore, the results of W-Tuckey's further test showed that the vitomolt dose of 1.5 g / kg of feed and 3.0 g / kg of feed was not significantly different from the control treatment, but the vitomolt dose of 4.5 g / kg of feed was significantly different from the control. The three treatments of adding vitomolts in the feed were not significantly different from the daily growth rate of vaname shrimp during the study.

The results of this study indicated that the higher the dose of vitomoltt added to the feed, the higher the daily growth rate of vaname shrimp (Table 1). This shows that the addition of vitomolts as phytoecdysteroid plays an important role in the growth of vaname shrimp. As it is known that molting is an important initial phase for crabs and shrimp because by moulting they can experience growth. Soumoff & Skinner (1983), the ecdyson hormone is able to increase the moulting process in crustaceans where the stimulation process of the moulting hormone in the premolt phase has an impact on the rapid moulting period and the synchronization of the moulting period. The ecdyson hormone thus has a good impact on metabolism and the moulting period in crab larvae. Gunamalai et al. (2003) also stated that ecdysteroid is the main steroid hormone in arthropods that regulates physiological functions, such as growth, metamorphosis, and reproduction.

Burdette (1962) in Klein (2004) also states that ecdysteroid apart from being a molting hormone also plays a role in increasing protein formation through increased mRNA synthesis. Donalson et al. (1978) stated that the most prominent metabolic action of steroids is the activation of protein metabolism. Protein synthesis is the most basic growth process, without large-scale protein production, growth will not occur (Jobling et al., 2001). The results of this study also showed that even though the protein content in the diet was low, only about 30%, the presence of

This publication is licensed under Creative Commons Attribution CC BY.
http://dx.doi.org/10.29322/IJSRP.10.11.2020.p10778

ecdysteroid supplementation, the body's protein formation was still high. This is relevant to Burdette (1962 in Klein 2004) explanation that ecdysteroid plays a role in increasing protein synthesis. According to Turner & Bagnara (1976), hormones not only play a role in stimulating action but also inhibiting action. There is a negative feedback mechanism in hormonal action. High hormone concentrations in the circulation give an indication for cells to perform inhibition in order to maintain balance (homeostasis).

IV. Survival Rate

The average survival rate (SR) of vaname shrimp at the end of the study can be seen in Table 2.

Table 2. Average survival rate of vaname shrimp at the end of the study

Vitomolt dose (g/kg of feed)	SR (%)
(A) Control	48.33 ± 20.82^{a}
(B) 1.5	75.00 ± 5.00^{ab}
(C) 3.0	80.00 ± 5.00^{b}
(D) 4.5	76.67 ± 7.64^{ab}

Note: Different letters in the SR column show significant differences between treatments (P <0.05)

The results of analysis of variance (ANOVA) showed that the addition of vitomolts to pelleted feed had a significant effect (p <0.05) on the survival rate of vaname shrimp. Furthermore, the results of W-Tuckey's further test showed that the vitomoltt dose of 3.0 g / kg of feed was significantly different from the treatment without vitomoltt but not different from the dose of 1.5 g and 4.5 g / kg of feed against the survival rate of vaname shrimp seeds.
The role of vitomolt as a supplement in feed is also seen in the survival rate of vaname shrimp. It can be seen that the vitomolt dose of 3.0 g / kg of feed is the optimum dose and when the dose is increased to 4.5 g / kg of feed, the survival rate of vaname shrimp decreases (Table 2). This shows that excess vitomolt supplementation actually has a negative impact on the survival rate of vaname shrimp. The results of research by Wahyuningsih (2008) and Susanti (2009) suggest that the right dose of ecdysteroid will stimulate crustaceans to molt optimally, while low doses cannot stimulate molting, otherwise too high doses cause inhibition.

V. Conclusions and Suggestions

The results showed a good response of the test shrimp with the application of vitomolts in feed. Vitomolt dose 3.0 g / kg of feed is the best treatment compared to other treatments. Based on the results of this study, it is recommended that the addition of vitomolts of 30 mg / kg of vaname shrimp seed feed is recommended.

References

[1] Aslamyah S. 2000. Pertumbuhan dan kelangsungan hidup ikan betutu (Oxyeleotris marmorata Blkr) yang diberi hormon metyltestosteron pada pakan dengan kadar protein berbeda. Jurnal Peternakan Universitas Hasanuddin, 8(2): 56-69.

[2] Donalson E.M., U.H.M Fegerlund, D.A.Higgs & J.R.Mc- Brede. 1978. Hormonal enhancement of growth. In: Hoar, W.S., D.J. Randall & J.R. Bret (Eds.). Fish Physiology. Vol VIII. Academic Press, New York. Pp. 456-597

[3] Fujaya, Y., S. Aslamyah, Z. Usman. 2011. Respon Molting, Pertumbuhan, dan Mortalitas Kepiting Bakau (Scylla olivacea) yang Disuplementasi Vitomolt melalui Injeksi dan Pakan Buatan. Jurnal ILMU KELAUTAN Vol. 16 (4) 211-218. ISSN 0853-7291.

[4] Gunamalai, V., Kirubagaran, R., & Subramoniam, T. (2003). Sequestration of ecdisteroid hormone into the ovary of the mole crab, Emerita asitica (Milne Edwards). Currents Science, 85(4), 493-496.

[5] Haliman, R.W. dan A.S. Dian, 2005. Udang vaname (Litopenaeus vaname): pembudidayaan dan prospek pasar udang putih yang tahan penyakit. Penebar Swadaya, Jakarta.

[6] Haryati, E. Saade dan Zainuddin. 2009. Formulasi dan aplikasi pakan untuk induk dan pembesaran: Aplikasi pakan buatan untuk peningkatan kualitas induk udang windu lokal. Laporan Penelitian Hibah Kompetitif Penelitian Sesuai Prioritas Nasional.

[7] Jobling M., T. Boujard, & D. Houlihan. 2001. Food Intake in Fish. Blackwell Science Ltd, A Blackwell Publishing Company

[8] Klein, R. 2004. Phytoecdysteroids. J. the American Herbalists Guild. Fall/Winter: 18-28

[9] Rosas, C., G. Cuzon, G. Gaxiola, Y. Le Priol, C. Pascual, J. rossignyol, F. Contreras, A. Sanchez & A. Van Wormhoudt. 2001. Metabolism and growth of juveniles of Litopenaeus vaname: effect of salinity and dietary carbohydrate levels. J. Exp.Mar.Biol.Ecol. 259:1-22

[10] Soumoff, C., & Skinner, D.M. (1983). Ecdysteroid titers during the molt cycle of the blue crab resemble those of other crustacean. The Biological Bulletin, 165, 321-329.

[11] Susanti, H. 2009. Pengaruh dosis vitomolt dalam pakan kepiting bakau (Scylla olivaceous) terhadap Molting. Skripsi. Fakultas Ilmu Kelautan dan Perikanan. Universitas Hasanuddin. Makassar. 34 hal

[12] Turner, C.D. & J.T. Bagnara. 1976. Endokrinologi Umum. Jogyakarta: Airlangga University Press. 746 hal

[13] Wahyuningsih, S.A. 2008. Pengaruh Dosis Penyuntikan Vitomolt terhadap Molting Kepiting Bakau (Scylla olivaceous). Skripsi. Fakultas Ilmu Kelautan dan Perikanan Universitas Hasanuddin. Makassar. 31 hal

[14] Zainuddin, Aslamyah, S., & Nur, K. 2019. The Effect of Dosage Combination and Feeding Frequency on Growth and Survival Rate of Vannamei Shrimp Juveniles in Ponds. In IOP Conference Series: Earth and Environmental Science (Vol. 370, No. 1, p. 012033). IOP Publishing.

[15] Zainuddin, Aslamyah, S., Nur, K., Hadijah. 2020. Substitution of sweet potato flour and corn starch to the growth, survival rate, feed conversion ratio and body chemical composition of juvenile Litopenaeus vannamei. Aquaculture, Aquarium, Conservation & Legislation, 13(5), 2497-2508.

Authors

First Author – Zainuddin, Fisheries Department, Faculty of Marine and Fisheries Science, Universitas Hasanuddin, Makassar, Corresponding E-mail: zainuddinlatief@gmail.com
Second Author – Y. Fujaya, Fisheries Department, Faculty of Marine and Fisheries Science, Universitas Hasanuddin, Makassar
Third Author – M.I. Djawad, Fisheries Department, Faculty of Marine and Fisheries Science, Universitas Hasanuddin, Makassar
Fourth Author – S. Aslamyah, Fisheries Department, Faculty of Marine and Fisheries Science, Universitas Hasanuddin, Makassar
Fifth Author – K. Nur, Fisheries Department, Faculty of Marine and Fisheries Science, Universitas Hasanuddin, Makassar
Fifth Author – Hadijah, Aquaculture Study Program, Faculty of Agriculture, Universitas Bosowa, Makassar

This publication is licensed under Creative Commons Attribution CC BY.

Effect of Vitomolt supplements in feed on growth and survival rate of white shrimp (Litopenaeus vannamei) seeds.

ORIGINALITY REPORT

22%	16%	18%	11%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	Z Zainuddin, S Aslamyah, K Nur, Hadijah. "The Effect of Dosage Combination and Feeding Frequency on Growth and Survival Rate of Vannamei Shrimp Juveniles in Ponds", IOP Conference Series: Earth and Environmental Science, 2019 Publication	5%
2	Submitted to Universitas Negeri Manado Student Paper	4%
3	repository.unhas.ac.id Internet Source	3%
4	Zainuddin, S Aslamyah, Hadijah. " Effect of carbohydrate types on water quality, proximate composition and glycogen content of white shrimp juvenile (,) ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	2%
5	Ali Osman, May Bin-Jumah, Mohamed E. Abd El-Hack, Ghada Elaraby et al. "Dietary	2%

supplementation of soybean glycinin can alter the growth, carcass traits, blood biochemical indices, and meat quality of broilers", Poultry Science, 2020
Publication

6	usnsj.com Internet Source	2%
7	www.ijsrp.org Internet Source	2%
8	aquasiana.org Internet Source	1%
9	www.plantphysiol.org Internet Source	1%
10	f1000researchdata.s3.amazonaws.com Internet Source	1%

Exclude quotes	On	Exclude matches	< 25 words
Exclude bibliography	On		

Effect of Vitomolt supplements in feed on growth and survival rate of white shrimp (Litopenaeus vannamei) seeds.

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4